

### **Minocyclin**

**Indikation:** Schwere Form der Akne. **Lyme-Borreliose**

**Kontraindikation:** Kinder unter 8 Jahren, Schwangerschaft, Stillzeit.

Beipackzettel beachten!

### **Doxycyclin**

**Indikation:** Bakterielle Infektionen, Gonorrhoe, Syphilis, **Lyme-Borreliose**. Infektionen im HNO-Bereich, Pneumonien durch Mykoplasmen, Rickettsien, Chlamydien. Chlamydien-Konjunktivitis und Trachom. Harnwegsinfekte (nur bei nachgewiesener Empfindlichkeit der Erreger), Prostatitis, Gonorrhoe, Syphilis, Cholera, Yersinien und Campylobacter – Infektionen, Shigellen bei Nachweis der Empfindlichkeit. Brucellose, Ornithose, Bartonellose, Listeriose, Rickettsiose, Meliodose, Pest, Granuloma inguinale. Ambulante Therapie von Gallenwegs-Infektionen. Hauterkrankungen, Akne vulgaris, Rosacea. Tropische Sprue. Morbus Whipple.

**Kontraindikation:** Kinder unter 8 Jahren, Schwangerschaft, Stillzeit.

Beipackzettel beachten!

### **Minocycline**

Indication: Severe form of acne. **Lyme Disease**

**Contraindications:** Children under 8 years of age, pregnancy, lactation.

Package inserts note!

### **Doxycycline**

**Indications:** Bacterial infections, gonorrhoea, syphilis, Lyme disease. Infections in the ENT field, pneumonia caused by mycoplasma, rickettsia, chlamydia. Chlamydia conjunctivitis and trachoma. Urinary tract infections (only in case of proven sensitivity of the pathogen), prostatitis, gonorrhoea, syphilis, cholera, Yersinia and Campylobacter - infections, shigella in detection sensitivity. Brucellosis, ornithosis, bartonellosis, listeriosis, rickettsiosis, Meliodose, plague, granuloma inguinale. Outpatient treatment of biliary tract infections. Skin disorders, acne, rosacea. Tropical sprue. Whipple's disease.

**Contraindications:** Children under 8 years of age, pregnancy, lactation.

Package inserts note!

Andersson H, Alestig K (1976) The penetration of doxycycline into CSF. Scand J Infect Dis Suppl. (9), 17-9.

Dattwyler RJ, Halperin JJ (1987) [Failure of tetracycline therapy in early Lyme disease.](#) Arthritis Rheum. 30(4), 448-50.

Dotevall L, Hagberg L (1989) Penetration of doxycycline into cerebrospinal fluid in patients treated for suspected Lyme neuroborreliosis. Antimicrob Agents Chemother 33(7), 1078-80

Cellerin L, Canfrere I, Ordroneau J, et al. (1994) **Acute eosinophilic pneumopathy induced by minocycline.** Rev Pneumol Clin 50(6), 325-8. [Abstract](#)

Piperno D, Donné C, Loire R, et al. (1995) Bronchiolitis obliterans organizing pneumonia associated with minocycline therapy: **a possible cause.** Eur Respir J. 8(6), 1018-20. <http://www.ncbi.nlm.nih.gov/pubmed/7589364>

Karlsson M, Hammers S, Nilsson-Ehle I, et al. (1996) Concentrations of doxycycline and penicillin G in sera and cerebrospinal fluid of patients treated for neuroborreliosis. *Antimicrob Agents Chemother* 40(5), 1104-7. [Abstract](#)

Ziska MH, Donta ST, Demarest FC (1996) Physician preferences in the diagnosis and treatment of Lyme disease in the United States. *Infection* 24(2), 182-6. [Abstract](#)

Donta ST (1997) Tetracycline therapy for chronic Lyme disease. *Clin Infect Dis* S52-6. [Abstract](#)

Steere AC (1997) Diagnosis and treatment of Lyme arthritis. *Med Clin North Am* 81(1), 179-94. [Abstract](#)

Shapiro LE, Knowles SR, Shear NH (1997) Comparative safety of tetracycline, minocycline, and doxycycline. *Arch Dermatol.* 133(10), 1224-30 <http://www.ncbi.nlm.nih.gov/pubmed/9382560>

Horowitz HW, Wormser GP (1998) Doxycycline revisited: an old medicine for emerging diseases. *Arch Intern Med* 158(2), 192-3. [Full Citation](#) | [Publisher Full Text](#)

Aberer E, Kehldorfer M, Binder B, et al. (1999) The outcome of Lyme borreliosis in children. *Wien Klin Wochenschr* 111(22-23), 941-4. [Abstract](#) | [Find Related Articles](#)

De Maria A, Primavera A (2000) Possibility of the use of oral long-acting tetracyclines in the treatment of Lyme neuroborreliosis. *Clin Infect Dis* 31(3), 848-9. [Full Citation](#) | [Publisher Full Text](#)

Bentas W, Karch H, Huppertz HI (2000) Lyme arthritis in children and adolescents: outcome 12 months after initiation of antibiotic therapy. *J Rheumatol* 27(8), 2025-30. [Abstract](#)

Dotevall L, Hagberg L (2000) Adverse effects of minocycline versus doxycycline in the treatment of Lyme neuroborreliosis. *Clin Infect Dis* 30(2), 410-1. [Full Citation](#) | [Publisher Full Text](#)

Cunha BA (2000) Minocycline versus doxycycline in the treatment of Lyme neuroborreliosis. *Clin Infect Dis* 30(1), 237-8. [Full Citation](#) | [Publisher Full Text](#)

Donta ST (2001) Treatment of patients with persistent symptoms and a history of Lyme disease. *N Engl J Med* 345(19), 1424; author reply 1425. [Full Citation](#)

Bernier C, Dréno B (2001) Minocycline. *Ann Dermatol Venereol* 128 (5), 627-37. [Abstract](#) | [Publisher Full Text](#)

Karkkonen K, Stiernstedt SH, Karlsson M (2001) Follow-up of patients treated with oral doxycycline for Lyme neuroborreliosis. *Scand J Infect Dis* 33(4), 59-62. [Abstract](#)

Bell CD, Kovacs K, Horvath E, et al. (2001) **Histologic, immunohistochemical, and ultrastructural findings in a case of minocycline-associated "black thyroid"**. *Endocr Pathol* 12(4), 443-51. [Abstract](#)

Lawson TM, Amos N, Bulgen B et al. (2001) **Minocycline-induced lupus: clinical features and response to rechallenge**. *Rheumatology* (2001) 40 (3): 329-335. doi: 10.1093/rheumatology/40.3.329 <http://rheumatology.oxfordjournals.org/content/40/3/329.full>

Bonnetblanc JM (2002) Doxycycline. *Ann Dermatol Venereol* 129(6-7), 874-82. [Abstract](#) | [Publisher Full Text](#)

Stricker RB, Lautin A (2003) The Lyme Wars: time to listen. *Expert Opin Investig Drugs* 12(10), 1609-14. [Abstract](#)

Kroun M (2003) Overvejelser / resume af publikationer vedr. **DOXYCYCLIN til behandling af BORRELIOSE og EHRlichiose**. <http://kroun.ulmarweb.dk/doxycyclin-dosis.htm>

van Steensel MA (2004) **Why minocycline can cause systemic lupus - a hypothesis and suggestions for therapeutic interventions based on it**. *Med Hypotheses* 63(1), 31-4. [Abstract](#)

Metz LM, Zhang Y, Yeung M et al. (2004) **Minocycline Reduces Gadolinium-Enhancing Magnetic Resonance Imaging Lesions in Multiple Sclerosis**. American Neurological Association. Published by Wiley-Liss, Inc., through Wiley Subscription Services. <http://owndoc.com/pdf/MS-lesions-minocycline.pdf>

Fritzsche M (2005) Chronic Lyme borreliosis at the root of multiple sclerosis--is a cure with antibiotics attainable? *Med Hypotheses* 64(3), 438-48. [Abstract](#)

Ogrinc K, Logar M, Lotric-Furlan S, et al. (2006) Doxycycline versus ceftriaxone for the treatment of patients with chronic Lyme borreliosis. *Wien Klin Wochenschr* 118(21-22), 696-701. [Abstract](#) | [Publisher Full Text](#)

Donta ST (2007) Lyme disease guidelines--it's time to move forward. *Clin Infect Dis* 44(8), 1134-5; author reply 1137-9. [Full Citation](#) | [Publisher Full Text](#)

Wormser GP, Halperin JJ (2008) Oral doxycycline for neuroborreliosis. *Lancet Neurol* 7(8), 665-6. [Full Citation](#)

Miyaoka T (2008) Clinical potential of minocycline for schizophrenia. *CNS Neurol Disord Drug Targets*. 7(4), 376-81. <http://www.ncbi.nlm.nih.gov/pubmed/18991666>

Ljøstad U, Skogvoll E, Eikeland R, et al. (2008) Oral doxycycline versus intravenous ceftriaxone for European Lyme neuroborreliosis: a multicentre, non-inferiority, double-blind, randomised trial. *Lancet Neurol* 7(8), 690-5. [Abstract](#) | [Publisher Full Text](#)

Aupee O, Almeras D, Le Garlantezec P, et al. (2009) Doxycycline. *Med Trop (Mars)* 69(6), 556-8. [Abstract](#) | [Find Related Articles](#)

Bernardino AL, Kaushal D, Philipp MT (2009) The antibiotics doxycycline and minocycline inhibit the inflammatory responses to the Lyme disease spirochete *Borrelia burgdorferi*. *J Infect Dis* 199(9), 1379-88. [Abstract](#)

Schröder J, Kirch W (2009) Lyme borreliosis: How should erythema migrans be treated in pregnant women? *Dtsch Med Wochenschr* 134(46), 2354. [Full Citation](#) | [Publisher Full Text](#)

Gulati RK (2010) Doxycycline in children?--the unanswered question. *Pediatr Dermatol* 27(4), 419. [Full Citation](#)

Sapi E, Kaur N, Anyanwu S et al. (2011) **Evaluation of in-vitro antibiotic susceptibility of different morphological forms of *Borrelia burgdorferi***. *Infect Drug Resist.* 4, 97-113. Published online May 3, 2011. doi: [10.2147/IDR.S19201](https://doi.org/10.2147/IDR.S19201) PMID: PMC3132871 <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3132871/> <http://www.dovepress.com/evaluation-of-in-vitro-antibiotic-susceptibility-of-different-morpholo-peer-reviewed-article-IDR>  
« **Conclusion: Antibiotics have varying effects on the different morphological forms of**

**B. burgdorferi. Persistence of viable organisms in round body forms and biofilm-like colonies may explain treatment failure and persistent symptoms following antibiotic therapy of Lyme disease ».**

Katsunori T, Iwasaki H, Ikegaya S, Ueda T (2012) Minocycline modulates cytokine and chemokine production in lipopolysaccharide-stimulated THP-1 monocytic cells by inhibiting IκB kinase α/β phosphorylation. *Translational Research* -, 1–11

Majewski MA (2014) Current opinion on the safety and efficacy of doxycycline including parenteral administration – A review. *Pol Ann Med*.  
<http://dx.doi.org/10.1016/j.poamed.2013.12.001>  
<http://www.sciencedirect.com/science/article/pii/S1230801314000022>

Berghoff W (2014) **Doxycyclin (Liquorgängigkeit)**.  
[http://www.praxis-berghoff.de/dokumente/berghoff150714/Kapitel\\_23-4-a\\_Doxycyclin\\_Liquorgaengigkeit.pdf](http://www.praxis-berghoff.de/dokumente/berghoff150714/Kapitel_23-4-a_Doxycyclin_Liquorgaengigkeit.pdf)

Moullan N et al. (2015) **Tetracyclines disturb mitochondrial function across eukaryotic models: a call for caution in biomedical research**. *Cell Reports*,  
doi:10.1016/j.celrep.2015.02.034.  
<http://www.cell.com/cell-reports/pdfExtended/S2211-1247%2815%2900180-1>

Keijmel SP, van Kasteren ME, Blox WA et al. (2015) **Cutaneous hyperpigmentation induced by doxycycline: a case series**. *Neth J Med*. 73(1), 37-40.  
<http://www.ncbi.nlm.nih.gov/pubmed/26219940>

Bremell D, Trollfors B (2017) **Doxycyklin kan ges till barn i alla åldrar - Ingen risk för missfärgning av tänder eller emaljhypoplasi**. *Lakartidningen*. 114. pii: ERIC. [Article in Swedish] <https://www.ncbi.nlm.nih.gov/pubmed/28850157>

**„Doxycycline can be given to children without risk of staining of teeth In Sweden, several hundred children are treated for Lyme neuroborreliosis annually, the majority of which are treated with intravenous ceftriaxone. Older tetracycline class antibiotics can cause permanent staining of the teeth. For doxycycline this has never been shown. Three publications on children exposed to doxycycline from three months of age show no risk of staining of the teeth. Changing the recommended treatment for children with Lyme neuroborreliosis to oral doxycycline would markedly simplify treatment for children and parents and reduce healthcare costs.“**

Rosenblat JD, McIntyre RS (2017) **Efficacy and tolerability of minocycline for depression: A systematic review and meta-analysis of clinical trials**. *J Affect Disord*. 227, 219-225. doi: 10.1016/j.jad.2017.10.042.  
<https://www.ncbi.nlm.nih.gov/pubmed/29102836>

**“Overall, a large antidepressant effect was observed for minocycline compared to placebo with good tolerability. The current analysis provides a proof-of-concept for the antidepressant effects of minocycline and provides impetus for future larger RCTs as well as identification of subgroups more likely to benefit from this intervention.”**

Bernt - Dieter Huismans Letzte Revision November 2017 [www.Huismans.click](http://www.Huismans.click)  
Back to top: <http://www.kabilahsystems.de/minocyclin.pdf>

